
Notes on the Correct Nomenclature of *Silene stewartii* S. A. Ghazanfar (Caryophyllaceae)

Subir Ranjan Kundu

Senior Research Fellow, Botanical Survey of India, 1, Sudder Street, Calcutta - 700016, India

ABSTRACT. During the preparation of a checklist of endemic taxa of Caryophyllaceae on the Indian subcontinent, the need for a nomenclatural correction of the following taxon became evident. Based on types, protologues, and herbarium (CAL, DD) specimens, *Silene stewartii* (Edgeworth) Majumdar is morphologically distinct from *Silene stewartii* S. A. Ghazanfar. The nomenclature of *Silene stewartii* S. A. Ghazanfar is corrected by establishing the new name *Silene subodhii* S. R. Kundu.

A new species of *Silene*, *S. stewartii* S. A. Ghazanfar, was published in 1982 based on material reported from the Pirpanjal range in Kashmir, northwest Himalaya (Ghazanfar, 1982; Ghazanfar & Nasir, 1986). The 1982 publication overlooked a note on a new combination of *Silene stewartii* (Majumdar, 1974) based on specimens reported from Lahul in Kashmir and Chamba of Himachal Pradesh (Majumdar, 1974, 1993).

Silene stewartii (Edgeworth) Majumdar, in Bull. Bot. Surv. India 16: 155. 1974. *Lychnis stewartii* Edgeworth, in Fl. Brit. India 1: 224. 1874. *Silene chambensis* Majumdar, in J. Indian Bot. Soc. 42: 649. 1963(1964).

Diagnostic features of *Silene stewartii* (Edgeworth) Majumdar bear little resemblance to *Silene stewartii* S. A. Ghazanfar (see Table 1). Further-

more, both taxa are described based on different type material. Hence, the later homonym *Silene stewartii* S. A. Ghazanfar (1982) is illegitimate, as the specific epithet is unavailable by its use as *S. stewartii* (Edgeworth) Majumdar in 1974, according to Article 53.1 of the *Code* (Greuter et al., 1994). A new binomial is proposed here, following Articles 7.3 and 33.2 of the *Code*.

Silene subodhii S. R. Kundu, nom. nov. Replaced name: *Silene stewartii* S. A. Ghazanfar, in Pakistan J. Bot. 14(2): 115. 1982. Not Majumdar, in Bull. Bot. Surv. India 16: 155. 1974. TYPE: Kashmir. Pir Panjal range, Frasnag, ±9000, 4.8 1947, R. R. & I. D. Stewart 23244 (holotype, RAW; isotype, NY).

The new specific epithet is given in honor of the author's father, Mr. Subodh Ranjan Kundu.

Literature Cited

- Edgeworth, M. P. & J. D. Hooker. 1874. Caryophyllaceae. In J. D. Hooker (editor), The Flora of British India 1: 212–246. L. Reeve, London.
- Ghazanfar, S. A. 1982. Two new *Silene* L. species from N.W. Himalayas. Pakistan J. Bot. 14(2): 113–116.
- & Y. J. Nasir. 1986. Caryophyllaceae. In E. Nasir & S. I. Ali (editors), Flora of West Pakistan. Fasc. No. 174: 1–125. Pakistan Agric. Res. Council, Islamabad.
- Greuter, W., F. R. Barrie, H. M. Burdet, W. G. Chaloner, V. Demoulin, D. L. Hawksworth, P. M. Jørgensen, D. H.

Table 1. Comparative morphological features of *S. stewartii* and *S. subodhii*.

<i>Silene stewartii</i>	<i>Silene subodhii</i>
1. Slender, grass-like herb	1. Short, woody herb
2. Stems 15–25 cm long	2. Stems 7–15 cm long
3. Leaves narrow, linear to lanceolate, 25–55 × 1.5–2 mm	3. Leaves dimorphic: a. basal leaves (leaves on main trunk) spatulate or oblong-lanceolate, 25–35 × 7–10 mm b. apical leaves (leaves on branches): linear-lanceolate, 18–25 × 5–6 mm
4. Calyx oblong-clavate, 8–9 mm long	4. Calyx campanulate, 10–11 mm
5. Petals ca. 7 × 2 mm	5. Petals ca. 8 × 1 mm
6. No. of styles: 3	6. No. of styles: 5
7. Seeds tuberculate, ca. 1.3 mm	7. Seeds papillate, ca. 1 mm

-
- Nicholson, P. C. Silva, P. Trehane & J. McNeill. 1994. International Code of Botanical Nomenclature (Tokyo Code). *Regnum Veg.* 131.
- Majumdar, N. C. 1963. Nomenclatural transfers in the Indian Caryophyllaceae—Silenoideae. *J. Indian Bot. Soc.* 42(1-4): 646-652.
- . 1974. *Lychnis stewartii* Edgeworth—Its identity, description, and distribution. *Bull. Bot. Surv. India* 16(1-4): 153-155.
- . 1993. Caryophyllaceae [P. Daniel, family editor]. *In* B. D. Sharma (editor), *Flora of India II*: 505-595. Botanical Survey of India, Calcutta.